



Overview of Rapi-G

What is Rapi-G ?

Rapi-G is the advanced generation method of 99.99% pure Chlorine dioxide. It is a molecule consisting 1 chlorine atom and 2 oxygen atoms. Abbreviated to **CLO₂**

It has a molecular weight of 67.45.

It is a gas at normal temperatures and pressures.

It has a melting point of -59°C.

It has a boiling point of 11°C

It is yellowish / green and has an odour similar to that of chlorine.

It is denser than air and is water soluble at standard temperatures and pressures up to 2500 ppm.

It is explosive in air at concentrations > 10%

It is prohibited from all form of transport, it is normally generated at the point of application.

It will decompose in the presence of UV, high temperatures, and high alkalinity (> ph 12).

Chlorine dioxide is not another form of chlorine. We can draw an analogue to hydrogen and hydrogen cyanide, they are both gases, have the same first name, but completely different properties. So too with chlorine dioxide and chlorine, indeed one molecule does make a big difference.

Chlorine Dioxide is defined in the USA as having "no elemental free chlorine" hence it does not chlorinate. It is because of this fact and the amazing chemistry of chlorine dioxide that it is slowly becoming an important tool in disinfection and oxidation in the world to-day.

The physical and chemical properties of chlorine dioxide outline below will unravel its amazing capabilities.

Chlorine dioxide does not dissociate in water.

It stays as chlorine dioxide therefore its ability to operate as a disinfectant / sanitizer is independent of pH.

Chlorine dioxide is an oxidant with a low redox potential.

It has a redox potential of +0.96 mV compared to chlorine of +1.36 mV. There is no relationship between redox and disinfecting efficacy.

Chlorine dioxide has a few specific chemical reactions.

From this property a number of very interesting properties are derived:

It has a very low toxicity rating, indeed some formulations have GRAS status. It is generally regarded as a " no - irritant".

It is not corrosive as a pure chlorine dioxide solution.

Its reactions are selective hence as an oxidant reagent consumption is maximised in

Rapi-G

AN ADVANCE DISINFECTANT

the redox reaction not through side reactions.

Chlorine dioxide has a very high efficacy against vegetative cells, for example, bacteria; fungi yeasts and molds; viruses; algae; and protozoa. It has little to no effect on human, animal and fish cells. It has been shown to have high efficacy against molluscs and acracides with unconfirmed reports suggesting some action against nematodes.

From the above properties it is not surprising then to learn that "chlorine dioxide does not constitute a risk against the environment". The Alliance for Environmental Technology (AET), is a group of 19 North American chemical manufacturers and forest product companies, established to promote proven and practical technologies to raise the environmental awareness has indicated that the "environmental risks of a modern paper mill using chlorine dioxide are INSIGNIFICANT."

The low oxidation potential of chlorine dioxide means that it can penetrate biofilm and indeed chlorine dioxide has been proven as the MOST effective chemical against biofilm. This has now been recognized by numerous organisations eg. Institute of Food Technologists in their report entitled "Microbial Attachment and Biofilm formation-A Scientific Summary, July '94 Food Technology. It has been clinically demonstrated that the presence of biofilm is the critical step in the infection pathway of legionellae. A simple and elegant solution is available in chlorine dioxide to overcome the problems related to having biofilm in a system. In terms of legionella control the single biggest problem is the formation of cysts, in the biofilm. Only chlorine dioxide and ozone have the capability of inactivating cysts!. Pulse dosing of a disinfectant is about a 1000 times more effective for biofilm control than low level continuous dosing.

CAUTION, is advised when one is running disinfection/sanitising programme during which one is eroding away the biofilm---theory and practice are indeed different bed mates.

chlorine dioxide is a factor lower in dosage for the same efficacy against bacteria and fungi when compared against any other standard disinfectant like chlorine, iodine, bromine, hydrogen peroxide, quaternary ammonium compounds (QUATS), glutaldehyde, phenolic and peroxyacetic acid formulations.

finally, chlorine dioxide can be easily and accurately measured in the food plant, potable water plant and for environmental applications. No other disinfectant / oxidiser can make this claim hence chlorine dioxide can easily meet GMP, HACCP, SQF or any other quality food safety management system or environmental system for consistency of performance.

In conclusion, therefore we have a disinfectant / sanitizer which is an oxidant with few chemical reactions, no pH limitations, very low toxicity, worldwide approval for drinking water, very high efficacy against micro-organisms, has a strong and measurable residual. The product when applied at use concentration in water will not corrode equipment; will not produce an environment harmful to workers or consumers.

Truly a wonderful product but it is not a magic bullet and it cannot solve all problems. We have examined the properties of chlorine dioxide that make it close to being the "ideal" biocide, however, the fact that it is a gas which cannot be compressed without exploding seemingly reduces its availability to be used.

Chlorine Dioxide Timeline

1811 first discovered by Sir Humphrey Davey.

1944 First commercial application. Used as a Biocide / Taste and Odour Control agent in domestic water at Niagara Falls in the USA.

1977 Three thousand municipal water systems achieving biological control using chlorine dioxide.

1980's chlorine dioxide gradually replacing chlorine in many industries.

Pulp and Paper industry as a bleaching agent.

Industrial water treatment as a biocide and as an odour control agent.

Food processing as a sanitizer.



- 1990's increasing used for the secondary disinfection of potable water.
- 2001 As the principal agent used in the decontamination of buildings in the United States after the anthrax attacks.
- 2005 Used after Hurricane Katrina to eradicate dangerous mold from house inundated by water from massive flooding.
- 2009 Used to prevent against H1N1 flu.

Chlorine Dioxide: The “Ideal” Biocide

Chlorine dioxide is an extremely effective disinfectant, which rapidly kills bacteria, viruses, and Giardia, and is also effective against Cryptosporidium. ClO₂ also improves taste and odour, destroys sulfides, cyanides, and phenols, controls algae, and neutralizes iron and manganese ions. It is an effective biocide at concentrations as low as 0.1 ppm (parts per million) and over a wide pH range. It is 10 times more soluble in water than chlorine, even in cold water. Unlike iodine, chlorine dioxide has no adverse effects on thyroid function. Chlorine dioxide is widely used by municipal water treatment facilities.

The term “chlorine dioxide” is misleading because chlorine is not the active element. Chlorine dioxide is an oxidizing, not a chlorinating agent. ClO₂ penetrates the cell wall and reacts with amino acids and the cytoplasm within the cell, killing the microorganism. Then by-product of this reaction is chlorite, which is harmless to humans.

For the super performance characteristics, chlorine dioxide has been described as the “ideal” biocide. It is now included in many drinking water hygiene programs around the globe. Complete testing has confirmed the safety of chlorine dioxide. This includes

extensive studies by the Environmental Protection Agency (EPA) and World Health Organization (WHO) Chlorine dioxide has been recognized by World Health Organization (WHO) as the most effective A1 disinfecting reagent.

Its usage was approved by Food and Drug Administration (FDA) and Environment Protection Agency (EPA).

Its status is also seen in the Report of FAO Codex Alimentarius, Food additive details Chlorine Dioxide.

Chlorine dioxide is approved and recommended by EPA as an environmental friendly drinking water additive to replace chlorine.

Chlorine dioxide has been called the “ideal” biocide for a number of reasons:

- It works against a wide variety of bacteria, yeasts, viruses, fungi, protozoa, spores, mold, mildews, and other microbes.
- It exhibits rapid kill of target organisms, often in seconds.
- It is effective at low concentrations and over a wide pH range.
- It biodegrades in the environment
- Unlike chlorine, it does not generate harmful by-products.

Chlorine Dioxide Germicidal Spectrum

Below table of some of organisms that chlorine dioxide has been tested with. Chlorine dioxide has proven effective at eliminating a wide range of organisms.

Bacteria	Leuconostoc mesenteroides	Tuberculosis
Blakeslea trispora	Listeria innocua ATCC 33090	Vancomycin-resistant Enterococcus faecalis(VRE)
Bordetella bronchiseptica	Listeria monocytogenes F4248	Vibrio strain Da-2
Brucella suis	Listeria monocytogenes F5069	Vibrio strain Sr-3
Burkholderia mallei	Listeria monocytogenes LCDC-81-861	Yersinia enterocolitica
Burkholderia pseudomallei	Listeria monocytogenes LCDC-81-886	Yersinia pestis

Rapi-G

AN ADVANCE DISINFECTANT

Campylobacter jejuni	Listeria monocytogenes Scott A	Yersinia ruckerii ATCC 29473
Clostridium botulinum	Methicillin-resistant Staphylococcus aureus (MRSA)	Viruses
Corynebacterium bovis	Multiple Drug Resistant Salmonella typhimurium (MDRS)	Adenovirus Type 40
Coxiella burneti (Q-fever)	Mycobacterium bovis	Canine Parvovirus
E. coli ATCC 11229	Mycobacterium fortuitum	Coronavirus
E. coli ATCC 51739	Pediococcus acidilactici PH3	Feline Calici Virus
E. coli K12	Pseudomonas aeruginosa	Foot and Mouth disease
E. coli O157:H7 13B88	Pseudomonas aeruginosa	Hantavirus
E. coli O157:H7 204P	Salmonella	Hepatitis A Virus
E. coli O157:H7 ATCC 43895	Salmonella spp.	Hepatitis B Virus
E. coli O157:H7 EDL933	Salmonella Agona	Hepatitis C Virus
E. coli O157:H7 G5303	Salmonella Anatum Group E	Algae/Fungi/Mold/Yeast

E. coli O157:H7 C7927	Salmonella Choleraesins ATCC 13076	Alternaria alternata
-----------------------	------------------------------------	----------------------

Erwinia carotovora (soft rot) Franscicella tularensis Fusarium sambucinum (dry rot)	Salmonella choleraesuis Salmonella Enterica (PT30) BAA-1045 Salmonella Enterica S. Enteritidis	Aspergillus aeneus Aspergillus aurolatus Aspergillus brunneo-uniseriatus
Fusarium solani var. coeruleum (dry rot) Helicobacter pylori	Salmonella Enterica S. Javiana Salmonella Enterica S. Montevideo	Aspergillus caespitosus Aspergillus cervinus
Helminthosporium solani (silver scurf) Klebsiella pneumonia	Salmonella Enteritidis E190-88 Salmonella Javiana	Aspergillus clavatonanicus Aspergillus clavatus
Lactobacillus acidophilus NRRL B1910 Lactobacillus brevis Lactobacillus buchneri Lactobacillus plantarum Legionella	Salmonella newport Salmonella Typhimurium C133117 Salmonella Anatum Group E Shigella Staphylococcus aureus	Aspergillus egyptiacus Aspergillus elongatus Aspergillus fischeri Aspergillus fumigatus Aspergillus giganteus
Legionella pneumophila Leuconostoc citreum TPB85	Staphylococcus aureus ATCC 25923 Staphylococcus faecalis ATCC 344	Aspergillus longivesica Aspergillus niger
Aspergillus ochraceus	Mucor flavus	Bacillus megaterium

Rapi-G

AN ADVANCE DISINFECTANT

Aspergillus parvathecius	Mucor indicus	Geobacillus stearothermophilus ATCC 12980/7953
Aspergillus sydowii	Mucor mucedo	Geobacillus stearothermophilus VHP
Aspergillus unguis	Mucor rademosus	Bacillus thuringiensis
Aspergillus ustus	Mucor ramosissimus	Chemical Decontamination
Aspergillus versicolor	Mucor saturnus	Mustard Gas
Botrytis species	Penicillium chrysogenum	Ricin Toxin
Candida spp.	Penicillium digitatum	dihydronicotinamide adenine dinucleotide
Candida albicans	Penicillium herquei	microcystin-LR (MC-LR)
Candida dubliniensis	Penicillium spp.	cylindrospermopsin (CYN)
Candida maltose	Phormidium boneri	Beta Lactams
Candida parapsilosis	Pichia pastoris	Amoxicillin
Candida sake	Poitrasia circinans	Ampicillin
Candida sojae	Rhizopus oryzae	Cefadroxil
Candida spp.	Saccharomyces cerevisiae	Cefazolin
Candida tropicalis	Stachybotrys chartarum	Cephalexin
Candida viswanathil	T-mentag (athlete's foot fungus)	Imipenem
Chaetomium globosum	Bacterial Spores	Penicillin G
Cladosporium cladosporioides	Alicyclobacillus acidoterrestris	Penicillin V
Debaryomyces etchellsii	Bacillus coagulans	Protozoa
Eurotium spp.	Bacillus anthracis	Cryptosporidium parvum Oocysts
Fusarium solani	Bacillus anthracis Ames	Chironomid larvae
Lodderomyces elongisporus	Bacillus atrophaeus	Microsporidia
Mucor circinelloides	Bacillus atrophaeus ATCC 49337	Encephalitozoon intestinalis

Disinfection By-Products of Chlorine Dioxide

The disinfection by-products (DBPs) of chlorine dioxide reactions are chlorite (ClO^-) and chlorate (ClO_2^-) and eventually chloride (Cl^-). The fate of any DBPs depends largely on the conditions at the time, concentration, temperature and the presence of other molecules.

Generally, it is the concentration of chlorite residuals that is the "monitored" DPB of chlorine dioxide. Modern generation systems are able to monitor the downstream residual DBP and adjust the dose rate to ensure that environmental limits are not breached. In special cases, downstream reactions can be used to remove excess chlorite residual from the water stream.

It is important to note that the DPBs of chlorine dioxide are easily managed with the correct experience and advice, and do not present nearly the same scale of problems as found with other biocide. Unlike ozone, chlorine dioxide does not oxidise bromide ion (Br^-) to bromate ion (BrO_3^-). Additionally, chlorine dioxide does not produce large amounts of aldehydes, ketones, or other DBPs that originate from the ozonisation of organic substances.

Approvals and Registrations for the use of Chlorine Dioxide USA Environment Protection Agency (EPA)

EPA approval for disinfectant / sanitizer with applications in food processing plants.

Rapi-G

AN ADVANCE DISINFECTANT

EPA approval for disinfectant of environmental surfaces such as floor, walls and ceiling in food processing plants, such as poultry, fish, meat, and in restaurants, dairies, bottling plants and breweries.

EPA approval for a terminal sanitising rinse for food contact surface in food processing plants, such as poultry, fish, meat, and in restaurants, dairies, bottling plants and breweries.

EPA approval for a sanitising rinse of uncut, unpeeled fruits and vegetables, at 5 ppm followed by a potable tap water rinse.

EPA approval for bacteriostatic in ice making plants and machinery.

EPA approval for treatment of stored potable water, at 5 ppm, for drinking water.

EPA bactericidal and fungicidal approval for hard non-porous surfaces in hospitals laboratories and medical environments.

EPA bactericidal and fungicidal approval for instruments in hospital and dental environment (Pending).

EPA bactericidal approval as a dental pumice disinfectant.

EPA approval for general disinfectant and deodorization of animal confinement

building, such as poultry, swine, barns and kennels.

EPA approval for the disinfection and deodorisation of ventilation systems and air conditioning duct work.

Food and Drug Administration (FDA)

FDA approval as a terminal sanitising rinse, not requiring a water rinse, on all food contact surface.

United States Department of Agriculture (USDA)

P-1 approval for bacterial and mould control in federally inspected meat and poultry processing plants for environmental surfaces.

D-2 approval as terminal sanitising rinse not requiring a water flush, on all food contact surfaces in food processing plant.

3-D approval for washing fruits and vegetables that are used as ingredients of meat, poultry and rabbit products by a potable water rinse.

G-5 approval for cooling and retort water treatment.

EU Codex Alimentarius

For use as an anti-microbial for incidental contact on food or surfaces that the food comes into contact with.

UK Government

Approved by the UK Secretary of State for the Environment under Regulation 25 (1)(a) of the Water Supply (Water Quality) (Amendments) Regulations 1991 (also in Scotland).

Approved as a disinfectant for service reservoirs, distribution, mains and waterworks apparatus.

Approved as a disinfectant and taste and odour control product for use in water that is supplied for drinking, washing, cooling and food production purposes on condition that the combined concentration of chlorine dioxide, chlorite and chlorate does not exceed 0.5 ppm entering supply.

Approved as a disinfectant by the Minister of Agriculture, Fisheries and Food and the Secretaries of State for Scotland and Wales for the Purposes of the Diseases of Animals (Approved Disinfectant) Order 1978 (As Amended) with corresponding approvals in Northern Ireland and Eire.

Approved for the Control of legionellae.

Approved by HS (G)70 for "The Control of legionellae including Legionnaires Disease" and MISC 150 the Technical Supplement to HS(G)70 "The Control of legionellae in Hot and Cold Water Systems".

Modes of Action of Chlorine Dioxide

Micro biocide Action

Rapi-G

AN ADVANCE DISINFECTANT

Chlorine dioxide is a stronger disinfectant than chlorine and chloramine. Ozone has great micro biocide effects, but limited residual disinfection capability. Recent research in the United States and Canada demonstrates that chlorine dioxide destroys enteroviruses, E. coli, amoebae and is effective against cryptosporidium cysts (Finch et al., 1997).

Chlorine dioxide exists in the water as ClO₂ (little or no dissociation) and thus is able to permeate through bacterial cell membranes and destroy bacterial cells (Junli et. Al, 1977b). its action on viruses involves adsorbing onto and penetrating the protein coat of the viral capsid and reacting with the viral RNA

(An RNA virus is a virus that has RNA (ribonucleic acid) as its genetic material. This nucleic acid is usually single-stranded RNA (ssRNA), but may be double-stranded RNA (dsRNA). Notable human diseases caused by RNA viruses include SARS, influenza, hepatitis C, West Nile fever, polio and measles.

The ICTV classifies RNA viruses as those that belong to Group III, Group IV or Group V of the Baltimore classification system of classifying viruses, and does not consider viruses with DNA intermediates in their life cycle as RNA viruses. Viruses with RNA as their genetic material but which include DNA intermediates in their replication cycle are called retroviruses, and comprise Group VI of the Baltimore classification. Notable human retroviruses include HIV-1 and HIV-2, the cause of the disease AIDS.


Another term for RNA viruses that explicitly excludes retroviruses is ribovirus.)

As a result, the genetic capability of the virus is damaged (Junli et. Al, 1977a). In comparison to chlorine, chlorine dioxide can be more effective as a disinfectant due to the fact that chlorine exists in the water as HOCl or OCl⁻. As a result, bacterial cell walls are negatively charged and repel these compounds, leading to less penetration and absorption of the disinfectant into the membranes.

Oxidant Action

The oxidant action of chlorine dioxide often improves the taste, odour, and colour of water. Chlorine dioxide reacts with phenolic compounds, humic substance, organics, and metal ions in the water.

For example, iron is oxidized by chlorine dioxide so that it precipitates out of the water in the form of iron hydroxide. The precipitate is then easily removed by filtration.



Chlorine dioxide reacts with organics, typically by oxidation reactions, and forms few chlorinated organic compounds. Free chlorine, in the presence of organic precursors can form trihalomethanes (THM's) and other halogenated compounds (Aieta and Berg, 1986).

Phenolic compounds presents in drinking water are due mainly to contamination from industrial sources. Such molecules, even when present in concentrations of micrograms per litre, give an unpleasant odour and taste. Chlorine dioxide reacts rapidly with phenols. This reaction may vary in different systems.

- (1) The formation of quinones or chloroquinones
- (2) The breaking of the aromatic ring and the formation of aliphatic derivatives.

Humic acid, a THM precursor, is oxidized by chlorine dioxide thus minimizing halogenated compound formation in secondary treatment (Aieta and Berg, 1986)

Function of Chlorine Dioxide

Chlorine dioxide can be used as a Disinfectant, Sanitizer, Tuberculocide, Virucide, Fungicide, Algaecide, Slimicide, and Deodorizer.

Rapi-G

AN ADVANCE DISINFECTANT

Chlorine dioxide is a powerful oxidising biocide and has been successfully used as a water treatment disinfectant for several decades in many countries. Rapid progress has been made in the technology for generation of the product and knowledge of its reactivity has increased with improved analytical techniques. Chlorine dioxide is a relatively stable radical molecule. It is highly soluble in water, has a boiling point of 110°C, absorbs light and breaks down into ClO^- and Cl^- . Because of its oxidising

properties chlorine dioxide acts on Fe^{2+} , Mn^{2+} and NO_3^- but does not act on Cl^- , NH_4^+ and Br^- when not exposed to light. These ions are generally part of the chemical composition of natural water.

Because of its radical structure, chlorine dioxide has a particular reactivity - totally different from that of chlorine or ozone. The latter behave as electron acceptors or are electrophilic, while chlorine dioxide has a free electron for a homopolar bond based on one of its oxygen. The electrophilic nature of chlorine or hypochloric acid can lead, through reaction of addition or substitution, to the formation of organic species while the radical reactivity of chlorine dioxide mainly results in oxycarbonyls. Generally chlorine dioxide rapidly oxidises phenol type compounds, secondary and tertiary amines, organic sulphides and certain hydrocarbon polycyclic aromatics such as benzopyrene, anthracene and benzoanthracene. The reaction is slow or non-existent on double carbon bonds, aromatic cores, quinionic and carboxylic structures as well as primary amines and urea.

The oxidising properties and the radical nature of chlorine dioxide make it an excellent virucidal and bactericidal agent in a large pH range. The most probable explanation is that in the alkaline media the permeability of living cell walls to gaseous chlorine dioxide radicals seems to be increased allowing an easier access to vital molecules. The reaction of chlorine dioxide with vital amino acids is one of the dominant processes of its action on bacteria and viruses.

Chlorine dioxide is efficient against viruses, bacteria and protozoan clumps usually found in raw water. A rise in pH level further increase its action against f2 bacteriophages, amoebic clumps, polioviruses and anterovirus. It is efficient against Giardia and has an excellent biocide effect against Cryptosporidia which are resistant to chlorine and chloramines. It has been demonstrated that ClO_2 has greater persistence than chlorine. In a recent report for dosages 3 times lower than those of chlorine at the station outlet, the residual of ClO_2 used alone was always higher than that of Cl_2 which also required 3 extra injections of chlorine in the distribution system.

The reduction of bad tastes and odour with ClO_2 is the result of the elimination of algae and on the negligible formation of organo-chlorinated derivatives. The latter formed under chlorination give rise to very unpleasant odours. By its action on dissolved organic materials, without the formation of organic halogen compounds, ClO_2 limits problems of taste and colour. In addition the low dosages used in post disinfection are an advantage. When chlorine dioxide replaces chlorine in a system it may take up to 15 days for the benefits of the change to become apparent. Changes should be made gradually to avoid problems of a sudden release of slime into the system.

Markets & Applications

Chlorine dioxide has a wide range of applications including:

Human Water Systems

- Treatment of Potable Water for Human Consumption
- Water Storage Systems Aboard Aircraft, Boats, RV's and Off-Shore Oil Rigs
- Municipal Well Waters
- Swimming Pools & Spas

Industrial

- Industrial water treatment
- Cooling and process water microbiological control
- Wastewater disinfection
- Cooling Towers

Rapi-G

AN ADVANCE DISINFECTANT

- Treatment of Ventilation Systems
- Mollusk control
- Odors control
- Iron and manganese removal
- Phenol oxidation
- Cyanide destruction
- Paper & pulp
- Influent Water Disinfection
- Backup on generators failures
- White water slimicide
- Iron Control
- Bleaching of specialty papers
- Oil & gas
- Microbiological control of oil wells and bores
- Sulfide destruction
- Pipeline and tank cleaning
- THM control
- Taste and odor control

Public Places

- Hospitals
- Microbiological control
- Lower risk of MRSA
- Cleaning
- Legionella prevention and control
- Hotels & leisure centers
- Disinfection of water system
- Biofilm removal in water system

Agricultural

- Horticulture
- Disinfection of irrigation water
- Cleaning of irrigation system
- Treatment of Agricultural Storage Facilities
- Treatment of Horticulture Work Area and Benches
- Treatment of Horticulture Pots and Flats
- Treatment of Horticulture Cutting Tools
- Treatment of Horticulture Bulbs
- Treatment of Greenhouse Glass, Walkways and Under Benches
- Treatment of Evaporative Coolers
- Treatment of Retention Basins and Ponds
- Treatment of Decorative Pools, Fountains and Water Displays
- Vegetables & fruit washing/processing

Food Processing

- Sanitizing Food Contact Surfaces
- Sanitizing Non-Food Contact Surfaces
- Sanitizing Food-Processing Equipment
- Ice Making Plants and Machinery
- Ice Manufacture
- Canning Retort and Pasteurizer Cooling Water

Rapi-G

AN ADVANCE DISINFECTANT

- Stainless Steel Transfer Lines, Hydro coolers and Pasteurizer
- Washing fruit and vegetables
- Washing fish and seafood
- Washing meat, poultry and processing equipment
- Extend shelf life and freshness of non-processed fruits and vegetables
- Process water for canned and frozen packaging
- Control of bacteria growth and bio fouling
- Control of salmonella and legionella
- Disinfection lines, holding tanks and other equipment
- Disinfect in beverage and water systems and lines
- Reduction of ammonia nitrogen concentration in recycled water
- Cleansing and rinsing of bottles
- Disinfect in beverage and water systems and lines
- CIP (Cleaning In Place)

Livestock

- Treatment of Drinking Water
- Disinfection of Animal Confinement Facilities
- Treatment of Animal Transport Vehicles
- Deodorization of Animal Holding Rooms, Sick Rooms and Work Rooms
- Control of Odor and Slime Forming Bacteria in Animal Confinement Facilities
- Disinfection of Poultry Chiller Water / Carcass Spray
- Treatment of Egg Room
- Treatment of Hatching Room
- Treatment of Incubator Room
- Treatment of Tray Washing Room and Loading Platform
- Treatment of Chick Room, Chick Grading Box and Sexing Room
- Hand Dip for Poultry Workers
- Shoe Bath Use

Aquaculture

- Live Fish Transport: Transport Water, Disease treatment during holding
- Disease prevention treatment
- Fish larval rearing
- Prawn larval rearing
- Spraying in feeds
- Treatment of diseases
- Fishing boats/Wholesale/Retail
- Dipping de-scaled and gutted fish
- Spray / Dipping on fish and prawns
- In sorting / grading water for prawns
- Ice manufacture
- Disinfection of display cabinet

Frequently Asked Questions of Chlorine Dioxide

Has chlorine dioxide been used before?

Chlorine dioxide has been recognized as an effective biocide for decades, and is used in a range of hygiene-related applications worldwide. Municipal water systems have used chlorine dioxide to treat drinking water for over 50 years.

Why couldn't I use chlorine dioxide before?

Prior to various chlorine dioxide delivery agent products, expensive mechanical generators or



relatively impure “stabilized” solutions were the only ways to make chlorine dioxide. The expense of capital equipment and the corrosiveness of the lower quality solutions prohibited the development of many horticultural applications.

Is chlorine dioxide safe?

The Niagara Falls New York water treatment plant first used chlorine dioxide for drinking water disinfection in 1944. Currently, there are approximately 400 – 500 water treatment plants in the United States and over 1000 in Europe utilizing ClO₂ to purify municipal drinking water systems. Numerous studies have shown chlorine dioxide, when used at the appropriate concentrations, has no adverse health effects, either by skin contact or ingestion.

Is chlorine dioxide toxic?

Fifty years of worker experience has demonstrated that ClO₂ is a safe compound when handled properly. World-wide, nearly 4.5 million pounds per day of chlorine dioxide are used in the production of pulp and paper. However, as with any and all disinfectant chemicals, if handled improperly, or consumed internally or absorbed or subjected to prolonged exposure, ClO₂ can be toxic. However, it is also this toxicity that makes ClO₂ a good water disinfectant agent.

Is chlorine dioxide environmentally friendly and does it create harmful by-products?

Chlorine dioxide is far more environmentally friendly than other oxidizing biocides and disinfectants including chlorine and bromine. Substituting chlorine dioxide for chlorine eliminates the formation of toxic halogenated disinfection by-products including trihalomethanes (THMs) and other chlorinated compounds that are harmful to the environment. In fact chlorine dioxide actually helps to remove substances that can form trihalomethanes. The disinfection is by oxidation as chlorine dioxide does not have either addition or substitution reactions associated with its chemistry.

What methods are used to detect chlorine dioxide?

Chlorine dioxide can be detected in several ways. Some of these methods such as DPD, Amperometric, and Iodometric are standardized, widely accepted and used.

Is Chlorine Dioxide expensive?

When compared to the cost of chlorine, the cost of ClO₂ is lower comparing efficiency and high range disinfection. However in those instances in which chlorine is not the preferred regulatory or environmental alternative, ClO₂ is a very attractive alternative. The costs are also less than that of other alternatives like ozone which can also be used for water treatment.

Can Chlorine Dioxide be stored safely?

No because explosive gas in the air (10%). Rapi-G provide a safety solution to produce “just in time”.

What legal provisions does chlorine dioxide carry?

Chlorine dioxide has a number of legal provisions by different states list in the follow table.

Time	State	Approved Bureau	Usage Range
1992		WHO	Drinking Water Disinfection
1985	U.S.A	FDA	Food Processing Equipment Sterilization
1985	EU	European Commission	Drinking Water Disinfection, food industry, medical, livestock husbandry, aquaculture, environment and public areas disinfection and sterilization
1987	Germany		Drinking Water Disinfection

Rapi-G

AN ADVANCE DISINFECTANT

1987	UK	Ministry of Health	Drinking Water Disinfection, hospital, livestock aquaculture, environment and public areas disinfection and sterilization
1987	U.S.A	EPA	Food processing plants, breweries, restaurants, environmental disinfection; Hospitals, labs and non-empty rigid surface equipment sterilization and removal mildew
1989	U.S.A	EPA	Storage water disinfection; Livestock, disinfection and deodorizing
1988	Japan	Ministry of Food Health	Drinking Water Disinfection
1987	Australia	Ministry of Health	No. 926 food Additives, food Bleacher
1987	China	Ministry of Health	Food industry, medical, pharmaceutical, livestock husbandry, aquaculture, environment and public areas disinfection and sterilization
1996	China	Ministry of Health	Food additives, fruits and vegetables Preservation
2002	U.S.A	FDA	Food processing equipment, pipe, crafts and arts equipment, especially in milk processing plant
2005	China	Ministry of Health	Drinking Water Disinfection
2011	Brazil	Ministry of Health	Drinking Water Disinfection, Food industry, Storage water disinfection, Livestock

Can chlorine dioxide be used in combination with other disinfectants?

Yes. Chlorine dioxide is often used in combination with chlorine in municipal drinking water plants in order to reduce the amount of trihalomethanes and HAAs that would be formed if chlorine were used alone. Chlorine dioxide is added as the primary disinfectant in order to remove a number of oxidisable compounds without forming chlorinated DBPs, while chlorine is added at low levels in order provide a residual biocide for use in the disinfection system.

Is chlorine dioxide different to chlorine?

Yes. While chlorine dioxide has chlorine in its name, its chemistry is radically different from that of chlorine. Chlorine dioxide is not "chlorine in disguise". Both chlorine dioxide and chlorine are oxidizing agents. They are electron receivers. Chlorine has the capacity to take in two electrons, whereas chlorine dioxide can absorb five. This is why chlorine dioxide is far more effective than chlorine as a disinfectant. Environmentally, chlorine dioxide is friendlier to the environment than chlorine. Chlorine dioxide does not form toxin trihalomethanes (THMs) or other chlorinated compounds that are harmful to the environment and associated with chlorine, sodium hypochlorite and hypochlorous acid.

What the difference between Rapi-G (chlorine dioxide) with other disinfectants?

Characters	Rapi-G	Chlorhexidine	Chlorine / Hypochlorite	Phenol	Aldehyde	NaOH	Alcohol
Resistance to Organic	Good	Ordinary	poor	General	good	good	General
Activity in Hard-water	Yes	Yes	Yes	No	Yes	Yes	Yes

Rapi-G

AN ADVANCE DISINFECTANT

Affect High Temperature	Result is best in 5-69 °C	No	Activity decreased below 40 °C	Activity increased	Result is best in 26-60 °C	No	No
PH Range	No effect	Alkaline	Acidic	Acidic	No effect	Alkaline	No effect
Anion Soap Compatibility	No	Yes	No	Yes	yes	Yes	No
Activity of Residue	No	Yes	No	Yes	No	Yes	No
Toxicity or Discomfort	No	No	Yes	Yes	Yes	Yes	Yes
Damage to Surface	No	No	Yes	No	Yes	Yes	No
Kill the Bacteria	Most	Part	Most	Most	Yes	Most	Most
Kill the Spores	Yes	Part	Part	No	Yes	Yes	No
Kill the Viruses	Yes	No	Part	Part	Yes	Yes	Part

How much is the Permissible Exposure Limit of chlorine dioxide?

The Occupational Safety and Health Administration (OSHA) has set safe exposure limits of 0.3 ppm (0.9 mg/m³) for 15 minutes and a time-weighted average of 0.1 ppm (0.3 mg/m³) for 8 hours of contact with chlorine dioxide gas.

What are advantages and disadvantages of chlorine dioxide?

Advantages:

- effective against a wide variety of bacteria, yeasts, viruses, fungi, protozoa, spores, molds, mildews, Cryptosporidium, algae and is more potent than chlorine over a short contact time

- destroys biofilms

- effective over wide pH (3.5 to 11)

- biodegradability in the environment

- prevents trihalomethanes (THM's) and bromate formation

- does not chlorinate organics

- readily dissolves in water and does not react with ammonia

- does not react with water to form free chlorine and hypochlorous acid

- does not react with water to form free chlorine

- is less corrosive than chlorine

- Selective oxidation reactions

- Cheaper than Ozone and more effective for odour, colour, bad taste, phenols reduction, iron and manganese reduction

Disadvantages:

- decomposes in sunlight

- must be generated on-site

Rapi-G

AN ADVANCE DISINFECTANT

EQUIPMENT EXPOSED TO Rapi-G (GASEOUS CHLORINE DIOXIDE) WITH NO ADVERSE EFFECTS	
Equipment	Results
Computers – Laptop and Desktop	No Effect
GC's and HPLC's	No Effect
VCR's / VHS Tapes	No Effect
Scales / Balances	No Effect
Microscopes	No Effect
Cameras / Camcorders	No Effect
MRI Machines	No Effect
X-Ray Machines	No Effect
Refrigerators / Freezers	No Effect
Incubators	No Effect
Autoclaves	No Effect
Cage / Rack Washers	No Effect
Temperature / Pressure / Rh Probes	No Effect
HEPA Filters	No Effect
Sterilizing Filters	No Effect
Solenoid Valves	No Effect

Biological Safety Cabinets	No Effect
Rigid Plastic / Stainless Steel / Flexible Wall Isolators	No Effect
Polypropylene	No Effect
HVAC ductwork	No Effect
Engines	No Effect
Siloxane Gel	No Effect
Ultrasound Machines	No Effect
Xenogen IVIS 2001 Imaging System	No Effect
Anesthesia Systems	No Effect
Ventilators	No Effect

Rapi-G

AN ADVANCE DISINFECTANT

Applications Of Rapi-G(Gas Chlorine Dioxide)



Isolators



**Pass-through
Rooms**



Surgical Suites



Large Animal Holding Rooms



Animal Holding Rooms



**Decontamination
Chamber**

Rapi-G

AN ADVANCE DISINFECTANT



Modified Autoclaves



Laboratories



Clean Rooms



Biological Safety Cabinets



Products and Components



HEPA Housings



Equipment Decontamination



Processing Rooms / Filing Lines



Necropsy Rooms



Buildings

Rapi-G

AN ADVANCE DISINFECTANT



Transport Van



BSL 1-4 Laboratories/Suites